

HEADS*UP

CONCUSSION IN HIGH SCHOOL SPORTS

A FACT SHEET FOR PARENTS

What is a concussion?

A concussion is a brain injury. Concussions are caused by a bump, blow, or jolt to the head or body. Even a “ding,” “getting your bell rung,” or what seems to be a mild bump or blow to the head can be serious.

What are the signs and symptoms?

You can’t see a concussion. Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days after the injury. If your teen reports **one or more** symptoms of concussion listed below, or if you notice the symptoms yourself, keep your teen out of play and seek medical attention right away.

Signs Observed by Parents or Guardians	Symptoms Reported by Athlete
<ul style="list-style-type: none"> • Appears dazed or stunned • Is confused about assignment or position • Forgets an instruction • Is unsure of game, score, or opponent • Moves clumsily • Answers questions slowly • Loses consciousness (even briefly) • Shows mood, behavior, or personality changes • Can’t recall events <i>prior</i> to hit or fall • Can’t recall events <i>after</i> hit or fall 	<ul style="list-style-type: none"> • Headache or “pressure” in head • Nausea or vomiting • Balance problems or dizziness • Double or blurry vision • Sensitivity to light or noise • Feeling sluggish, hazy, foggy, or groggy • Concentration or memory problems • Confusion • Just not “feeling right” or is “feeling down”

How can you help your teen prevent a concussion?

Every sport is different, but there are steps your teens can take to protect themselves from concussion and other injuries.

- Make sure they wear the right protective equipment for their activity. It should fit properly, be well maintained, and be worn consistently and correctly.

- Ensure that they follow their coaches’ rules for safety and the rules of the sport.
- Encourage them to practice good sportsmanship at all times.

What should you do if you think your teen has a concussion?

- 1. Keep your teen out of play.** If your teen has a concussion, her/his brain needs time to heal. Don’t let your teen return to play the day of the injury and until a health care professional, experienced in evaluating for concussion, says your teen is symptom-free and it’s OK to return to play. A repeat concussion that occurs before the brain recovers from the first—usually within a short period of time (hours, days, or weeks)—can slow recovery or increase the likelihood of having long-term problems. In rare cases, repeat concussions can result in edema (brain swelling), permanent brain damage, and even death.
- 2. Seek medical attention right away.** A health care professional experienced in evaluating for concussion will be able to decide how serious the concussion is and when it is safe for your teen to return to sports.
- 3. Teach your teen that it’s not smart to play with a concussion.** Rest is key after a concussion. Sometimes athletes wrongly believe that it shows strength and courage to play injured. Discourage others from pressuring injured athletes to play. Don’t let your teen convince you that s/he’s “just fine.”
- 4. Tell all of your teen’s coaches and the student’s school nurse about ANY concussion.** Coaches, school nurses, and other school staff should know if your teen has ever had a concussion. Your teen may need to limit activities while s/he is recovering from a concussion. Things such as studying, driving, working on a computer, playing video games, or exercising may cause concussion symptoms to reappear or get worse. Talk to your health care professional, as well as your teen’s coaches, school nurse, and teachers. If needed, they can help adjust your teen’s school activities during her/his recovery.

If you think your teen has a concussion:

Don’t assess it yourself. Take him/her out of play. Seek the advice of a health care professional.

It’s better to miss one game than the whole season.

For more information and to order additional materials **free-of-charge**, visit: www.cdc.gov/Concussion.

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION



Suggested Return to Play (RTP) Procedures After a Concussion

1. Return to activity and play is a medical decision. The athlete must meet all of the following criteria in order to progress to activity:

Asymptomatic at rest and with exertion (including mental exertion in school) AND have written clearance from their primary care provider or concussion specialist (athlete must be cleared for progression to activity by a physician other than an Emergency Room physician, if diagnosed with a concussion).

2. Once the above criteria are met, the athlete will be progressed back to full activity following the step-wise process detailed below. (This progression must be closely supervised by a Certified Athletic Trainer. If your school does not have an athletic trainer, then the coach must have a very specific plan to follow as directed by the athlete's physician).
3. Progression is individualized, and will be determined on a case by case basis. Factors that may affect the rate of progression include: previous history of concussion, duration and type of symptoms, age of the athlete, and sport/activity in which the athlete participates. An athlete with a prior history of concussion, one who has had an extended duration of symptoms, or one who is participating in a collision or contact sport may progress more slowly. Further, concussion in females tends to take longer to heal and should be carefully watched.
4. Stepwise progression as described below:

Step 1: Complete cognitive rest. This may include staying home from school or limiting school hours (and studying) for several days. Activities requiring concentration and attention may worsen symptoms and delay recovery.

Step 2: Return to school full-time.

Step 3: Light exercise. This step cannot begin until the athlete is no longer having concussion symptoms and is cleared by a physician for further activity. At this point the athlete may begin walking or riding an exercise bike. No weight-lifting or jumping/bouncing up and down.

Step 4: Running in the gym or on the field. No helmet or other equipment.

Step 5: Non-contact training drills in full equipment. Weight-training can begin.

Step 6: Full contact practice or training.

Step 7: Play in game. Must be cleared by physician before returning to play.

- *The athlete should spend 1 to 2 days at each step before advancing to the next. If post-concussion symptoms occur at any step, the athlete must stop the activity and the treating physician must be contacted. Depending upon the specific type and severity of the symptoms, the athlete may be told to rest for 24 hours and then resume activity at a level one step below where he or she was at when the symptoms occurred.*